



Human papillomavirus quantification in urine and cervical samples by using the Mx4000 and LightCycler general real-time PCR systems

Submitted by Alexandra Ducancelle on Thu, 02/07/2019 - 15:44

Titre	Human papillomavirus quantification in urine and cervical samples by using the Mx4000 and LightCycler general real-time PCR systems
Type de publication	Article de revue
Auteur	Payan, Christopher [1], Ducancelle, Alexandra [2], Aboubaker, Mohamed H [3], Caer, Julien [4], Tapia, Malena [5], Chauvin, Amelie [6], Peyronnet, Damien [7], Le Hen, Elodie [8], Arab, Zohra [9], Legrand, Marie-Christine [10], Tran, Adissa [11], Postec, Edith [12], Tourmen, Françoise [13], Avenel, Martine [14], Malbois, Chantal [15], De Brux, Marie-Anne [16], Descamps, Philippe [17], Lunel-Fabiani, Françoise [18]
Editeur	American Society for Microbiology
Type	Article scientifique dans une revue à comité de lecture
Année	2007
Langue	Anglais
Date	Mars 2007
Numéro	3
Pagination	897-901
Volume	45
Titre de la revue	Journal of clinical microbiology
ISSN	0095-1137
Mots-clés	Cervix Uteri [19], DNA, Viral [20], Female [21], Humans [22], Mass Screening [23], Papillomaviridae [24], Papillomavirus Infections [25], Polymerase Chain Reaction [26], Sensitivity and Specificity [27], Urine [28], Uterine Cervical Neoplasms [29]

Résumé en anglais

During the last decade, growing efforts have focused on human papillomavirus (HPV) detection using liquid hybridization, conventional PCR, and real-time PCR-based methods to increase the overall proportion of patients participating in cervical cancer screening procedures. We proposed a new general HPV DNA real-time PCR on the Mx4000 (Stratagene) and LightCycler (Roche Diagnostics) systems usable for both cervical scrape specimens and urine samples. A linear range was obtained from 5 DNA copies to 8 log(10) DNA copies/ml, and intra- and interassay variations were between 1.8 and 4%. Cervical carcinoma and HPV DNA screening was performed in 333 individual women referred for gynecological examination at the university hospitals of Angers and Brest and enrolled in the PapU study. Among cervical specimens (n = 333), 45% were positive for HPV DNA, with a mean viral load at 5.00 log/ml (+/- 1.73). Among urine samples (n = 177), 37% were positive with a significant 50-fold-lower mean viral load (3.77 +/- 1.32 log/ml; P < 0.0001). Kappa agreement for HPV DNA between cervical and urine specimens was excellent (93%). Thus, we developed a highly sensitive and quantitative general HPV DNA real-time PCR method that allows mass screening of patients with HPV infection. The ongoing longitudinal and prospective multicenter PapU study should give us the opportunity to validate this method adapted to HPV DNA screening in urine samples in a larger population.

URL de la notice	http://okina.univ-angers.fr/publications/ua18810 [30]
DOI	10.1128/JCM.02022-06 [31]
Lien vers le document	https://jcm.asm.org/content/45/3/897 [32]
Titre abrégé	J. Clin. Microbiol.
Identifiant (ID) PubMed	17229868 [33]
PubMed Central ID	PMC1829135

Liens

- [1] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=5349>
- [2] <http://okina.univ-angers.fr/a.ducancelle/publications>
- [3] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33725>
- [4] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33726>
- [5] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33727>
- [6] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33728>
- [7] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=20639>
- [8] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33729>
- [9] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33730>
- [10] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33731>
- [11] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33732>
- [12] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33733>
- [13] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33734>
- [14] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33735>
- [15] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33736>
- [16] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=33737>
- [17] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=16486>
- [18] <http://okina.univ-angers.fr/f.lunel/publications>
- [19] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=27093>
- [20] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=10217>

- [21] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=1075>
- [22] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=991>
- [23] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=15855>
- [24] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=10166>
- [25] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=10174>
- [26] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=8595>
- [27] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=7545>
- [28] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=16884>
- [29] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=27094>
- [30] <http://okina.univ-angers.fr/publications/ua18810>
- [31] <http://dx.doi.org/10.1128/JCM.02022-06>
- [32] <https://jcm.asm.org/content/45/3/897>
- [33] <http://www.ncbi.nlm.nih.gov/pubmed/17229868?dopt=Abstract>

Publié sur *Okina* (<http://okina.univ-angers.fr>)